

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
13 March 2003 (13.03.2003)

PCT

(10) International Publication Number
WO 03/020266 A1

(51) International Patent Classification⁷: **A61K 31/395**,
A61P 35/00, A61K 31/4706 // (A61K 31/395, 31:4706)

(21) International Application Number: PCT/US02/24841

(22) International Filing Date: 6 August 2002 (06.08.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/310,646 7 August 2001 (07.08.2001) US

(71) Applicant: **WYETH** [US/US]; Five Giralda Farms, Madison, NJ 07940-0874 (US).

(72) Inventors: **RABINDRAN**, Sridhar, Krishna; 2 Pamela Drive, Chestnut Ridge, NY 10977 (US). **GIBBONS**, James, J., Jr.; 33 Terrace Drive, Westwood, NJ 07675 (US).

(74) Agents: **MILOWSKY**, Arnold, S.; Wyeth, Patent Law Dept., Five Giralda Farms, Madison, NJ 07940-0874 et al. (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ANTINEOPLASTIC COMBINATIONS



(57) Abstract: This invention provides the use of a combination of CCI-779 and EKB-569 in the treatment of neoplasms.

WO 03/020266 A1

ANTINEOPLASTIC COMBINATIONS

BACKGROUND OF THE INVENTION

5 This invention relates to the use of combinations of rapamycin 42-ester with 3-hydroxy-2-(hydroxymethyl)-2-methylpropionic acid (CCI-779) and 4-dimethylamino-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-ethoxy-quinolin-6-yl]-amide (EKB-569).

10 Rapamycin is a macrocyclic triene antibiotic produced by Streptomyces hygroscopicus, which was found to have antifungal activity, particularly against Candida albicans, both in vitro and in vivo [C. Vezina et al., J. Antibiot. 28, 721 (1975); S.N. Sehgal et al., J. Antibiot. 28, 727 (1975); H. A. Baker et al., J. Antibiot. 31, 539 (1978); U.S. Patent 3,929,992; and U.S. Patent 3,993,749]. Additionally, rapamycin alone (U.S. Patent 4,885,171) or in combination with picibanil (U.S. Patent 15 4,401,653) has been shown to have antitumor activity.

The immunosuppressive effects of rapamycin have been disclosed in FASEB 3, 3411 (1989). Cyclosporin A and FK-506, other macrocyclic molecules, also have been shown to be effective as immunosuppressive agents, therefore useful in preventing transplant rejection [FASEB 3, 3411 (1989); FASEB 3, 5256 (1989); 20 R. Y. Calne et al., Lancet 1183 (1978); and U.S. Patent 5,100,899]. R. Martel et al. [Can. J. Physiol. Pharmacol. 55, 48 (1977)] disclosed that rapamycin is effective in the experimental allergic encephalomyelitis model, a model for multiple sclerosis; in the adjuvant arthritis model, a model for rheumatoid arthritis; and effectively inhibited the formation of IgE-like antibodies.

25 Rapamycin is also useful in preventing or treating systemic lupus erythematosus [U.S. Patent 5,078,999], pulmonary inflammation [U.S. Patent 5,080,899], insulin dependent diabetes mellitus [U.S. Patent 5,321,009], skin disorders, such as psoriasis [U.S. Patent 5,286,730], bowel disorders [U.S. Patent 5,286,731], smooth muscle cell proliferation and intimal thickening following vascular 30 injury [U.S. Patents 5,288,711 and 5,516,781], adult T-cell leukemia/lymphoma [European Patent Application 525,960 A1], ocular inflammation [U.S. Patent 5,387,589], malignant carcinomas [U.S. Patent 5,206,018], cardiac inflammatory disease [U.S. Patent 5,496,832], and anemia [U.S. Patent 5,561,138].

Rapamycin 42-ester with 3-hydroxy-2-(hydroxymethyl)-2-methylpropionic acid (CCI-779) is ester of rapamycin which has demonstrated significant inhibitory effects on tumor growth in both in vitro and in vivo models. The preparation and use of hydroxyesters of rapamycin, including CCI-779, are disclosed in U.S. Patent
5 5,362,718.

CCI-779 exhibits cytostatic, as opposed to cytotoxic properties, and may delay the time to progression of tumors or time to tumor recurrence. CCI-779 is considered to have a mechanism of action that is similar to that of sirolimus. CCI-779 binds to and forms a complex with the cytoplasmic protein FKBP, which inhibits an enzyme,
10 mTOR (mammalian target of rapamycin, also known as FKBP12-rapamycin associated protein [FRAP]). Inhibition of mTOR's kinase activity inhibits a variety of signal transduction pathways, including cytokine-stimulated cell proliferation, translation of mRNAs for several key proteins that regulate the G1 phase of the cell cycle, and IL-2-induced transcription, leading to inhibition of progression of the cell
15 cycle from G₁ to S. The mechanism of action of CCI-779 that results in the G₁ S phase block is novel for an anticancer drug.

In vitro, CCI-779 has been shown to inhibit the growth of a number of histologically diverse tumor cells. Central nervous system (CNS) cancer, leukemia (T-cell), breast cancer, prostate cancer, and melanoma lines were among the most
20 sensitive to CCI-779. The compound arrested cells in the G₁ phase of the cell cycle.

In vivo studies in nude mice have demonstrated that CCI-779 has activity against human tumor xenografts of diverse histological types. Gliomas were particularly sensitive to CCI-779 and the compound was active in an orthotopic glioma model in nude mice. Growth factor (platelet-derived)-induced stimulation of a human
25 glioblastoma cell line in vitro was markedly suppressed by CCI-779. The growth of several human pancreatic tumors in nude mice as well as one of two breast cancer lines studied in vivo also was inhibited by CCI-779.

Protein tyrosine kinases are a class of enzymes that catalyze the transfer of a
30 phosphate group from ATP or GTP to tyrosine residue located on protein substrates. Protein tyrosine kinases clearly play a role in normal cell growth. Many of the growth factor receptor proteins function as tyrosine kinases and it is by this process that they effect signaling. The interaction of growth factors with these receptors is a necessary event in normal regulation of cell growth. However, under certain conditions, as a

result of either mutation or overexpression, these receptors can become deregulated; the result of which is uncontrolled cell proliferation which can lead to tumor growth and ultimately to the disease known as cancer [Wilks A.F., *Adv. Cancer Res.*, 60, 43 (1993) and Parsons, J.T.; Parsons, S.J., *Important Advances in Oncology*, DeVita V.T. Ed., J.B. Lippincott Co., Phila., 3 (1993)]. Among the growth factor receptor kinases and their proto-oncogenes that have been identified and which are targets of the compounds of this invention are the epidermal growth factor receptor kinase (EGF-R kinase, the protein product of the erbB oncogene), and the product produced by the erbB-2 (also referred to as the neu or HER2) oncogene. Since the phosphorylation event is a necessary signal for cell division to occur and since overexpressed or mutated kinases have been associated with cancer, an inhibitor of this event, a protein tyrosine kinase inhibitor, will have therapeutic value for the treatment of cancer and other diseases characterized by uncontrolled or abnormal cell growth. For example, overexpression of the receptor kinase product of the erbB-2 oncogene has been associated with human breast and ovarian cancers [Slamon, D. J., et. al., *Science*, 244, 707 (1989) and *Science*, 235, 1146 (1987)]. Deregulation of EGF-R kinase has been associated with epidermoid tumors [Reiss, M., et. al., *Cancer Res.*, 51, 6254 (1991)], breast tumors [Macias, A., et. al., *Anticancer Res.*, 7, 459 (1987)], and tumors involving other major organs [Gullick, W.J., *Brit. Med. Bull.*, 47, 87 (1991)]. Because of the importance of the role played by deregulated receptor kinases in the pathogenesis of cancer, many recent studies have dealt with the development of specific PTK inhibitors as potential anti-cancer therapeutic agents [some recent reviews: Burke, T.R., *Drugs Future*, 17, 119 (1992) and Chang, C.J.; Geahlen, R.L., *J. Nat. Prod.*, 55, 1529 (1992)].

25

4-Dimethylamino-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-ethoxy-quinolin-6-yl]-amide (EKB-569) is an EGFR kinase inhibitor which has significant inhibitory effects on tumor growth in both in vitro and in vivo models. The preparation and use of EGFR kinase inhibitors, such as EKB-569, are disclosed in U.S. Patent 6,002,008.

30

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows cytotoxicity curves of EKB-569, CCI-779, and combinations of EKB-569 + CCI-779 in HCT116 cells.

FIG. 2 shows isobolograms (at the 50% effect level) of a EKB-569 + CCI-779 combination.

FIG. 3 shows isobolograms for EKB-569 + CCI-779 combinations derived from different endpoints ranging from 50-65%.

FIG. 4 shows a 3-dimensional analysis of the synergistic interaction of a EKB-569 + CCI-779 combination.

FIG. 5 shows a contour plot of the 3-dimensional synergy plot of a EKB-569 + CCI-779 combination.

DESCRIPTION OF THE INVENTION

This invention provides the use of combinations of CCI-779 and EKB-569 as antineoplastic combination chemotherapy. In particular, these combinations are useful in the treatment of renal cancer, soft tissue cancer, breast cancer, neuroendocrine tumor of the lung, cervical cancer, uterine cancer, head and neck cancer, glioma, non-small lung cell cancer, prostate cancer, pancreatic cancer, lymphoma, melanoma, small cell lung cancer, ovarian cancer, colon cancer, esophageal cancer, gastric cancer, leukemia, colorectal cancer, and unknown primary cancer. This invention also provides combinations of CCI-779 and EKB-569 for use as antineoplastic combination chemotherapy, in which the dosage of either CCI-779 or EKB-569 or both are used in subtherapeutically effective dosages.

As used in accordance with this invention, the term "treatment" means treating a mammal having a neoplastic disease by providing said mammal an effective amount of a combination of CCI-779 and EKB-569 with the purpose of inhibiting growth of the neoplasm in such mammal, eradication of the neoplasm, or palliation of the mammal.

As used in accordance with this invention, the term "providing," with respect to providing the combination, means either directly administering the combination, or administering a prodrug, derivative, or analog of one or both of the components of the combination which will form an effective amount of the combination within the body.

The preparation of CCI-779 is described in U.S. Patent 5,362,718, which is hereby incorporated by reference. An improved preparation of CCI-779 is disclosed in US patent application SN 09/670,358, which is hereby incorporated by reference. When CCI-779 is used as an antineoplastic agent, it is projected that initial i.v. infusion dosages will be between about 0.1 and 100 mg/m² when administered on a daily dosage regimen (daily for 5 days, every 2-3 weeks), and between about 0.1 and 1000 mg/m² when administered on a once weekly dosage regimen. Oral or intravenous infusion are the preferred routes of administration, with intravenous being more preferred.

10

EKB-569 can be prepared according to the procedures described in US Patent 6,002,008, which is incorporated by reference. Preferred procedures for the preparation of EKB-569 are provided herein. When EKB-569 is used as an antineoplastic agent it is projected that the initial oral dosage will be between 1 and 100 mg per day. Depending on patient tolerance, EKB-569 can be administered daily for a treatment period, such as 14 days, followed by a rest period (no drug administered), or can be administered on a continuous basis for a longer treatment period (for example, 6 months or longer).

20

The antineoplastic activity of the CCI-779 plus EKB-569 combination was confirmed in *in vitro* standard pharmacological test procedure; the following briefly describes the procedure used and the results obtained.

Cell Proliferation Procedure - HCT 116 colon adenocarcinoma cells were maintained in RPMI 1640 medium (Life Technologies, Inc., Gaithersburg, MD) supplemented with 10% fetal bovine serum (FBS, Life Technologies) and 50 µg/ml gentamicin (Life Technologies) under 7% CO₂ at 37°C. Cells were plated in 96-well microtiter dishes (6000 cells/well) in 200 µl RPMI 1640 medium containing 5% FBS and 50 µg/ml gentamicin and incubated overnight at 37°C. Compound dilutions were prepared in the same medium, at 5X final concentration, and 50 µl of the drug dilution was added to the cell-containing wells. For studies involving combinations of two drugs, serial dilutions of one compound were prepared in the presence of a fixed dose of a second compound. Alternatively, a checkerboard dilution series was employed. Cells were cultured for three days in the presence of the drugs. Untreated cells were

30

included as controls. The percentage of surviving cells was determined using sulforhodamine B (SRB, Sigma-Aldrich, St Louis, MO), a protein binding dye. Cellular protein was precipitated in each well by the addition of 50 μ l of 50% cold trichloroacetic acid. After 1 hour, the plates were washed extensively in water and dried. SRB dye reagent (0.4% SRB in 1% acetic acid, 80 μ l per well) was added and plates were kept at room temperature for ten minutes. Plates were then washed thoroughly in 1% acetic acid and dried. Cell-associated dye was dissolved in 10 mM Tris (150 μ l) and the absorbance was read at 540 nm in a microtiter plate reader. The concentration of compound that caused a fixed percentage inhibition of growth was determined by plotting cell survival (relative to untreated cells) against the compound dose.

Synergy Evaluation - Isobolograms were used to study the interaction of two pharmacological agents. Here, the concentration of each drug alone which produces a certain endpoint (e.g 50% inhibition of cell growth, IC_{50}), is plotted on the two graphical axes. The straight line connecting the two points represents equally effective concentrations of all combinations of the two drugs if the interaction is purely additive. A shift of the isobologram to the left of the predicted cytotoxicity (curve with concave side up) represents a synergistic interaction. Conversely, a shift to the right (isobologram with the convex side up) represents an antagonistic interaction. When isobolograms for different endpoints were plotted on the same graph, the concentration of each drug was expressed as the fraction of the concentration of each drug alone that produced the same effect. This produces a symmetrical isobologram with unit-less measures on each axis, and allows a direct comparison of different endpoints.

A second model for studying drug interactions was proposed by Prichard and Shipman [Antiviral Research. 14: 181-206 (1990)]. This is a 3-dimensional model: one for each drug and the third for the biological effect. Theoretical additive interactions are calculated from the individual dose-response curves, based on a dissimilar sites model of additivity (Bliss independence). The calculated additive surface, representing predicted cytotoxicity is subtracted from the experimental surface to reveal areas of enhanced toxicity (synergy) or reduced toxicity (antagonism). The resulting surface appears as a horizontal plane at 0% inhibition

above the calculated additive surface, if the interaction is additive. Peaks and valleys deviating from this plane are indicative of synergy and antagonism, respectively. MacSynergyII, a Microsoft Excel-based software was used to perform all calculations automatically. This spreadsheet calculates the theoretical additive interactions, and
5 locates and quantifies synergistic or antagonistic interactions that are significant at the 95% confidence levels. The results were plotted as a 3-dimensional plot, or as a contour plot.

Results - HCT 116 cells were chosen as they express low, but detectable
10 levels of EGFR, and are sensitive to inhibition by EGFR inhibitors. The cells are somewhat resistant to CCI-779, but are inhibited by high doses (5-10 $\mu\text{g/ml}$) of this drug. HCT-116 cells were cultured in the presence of EKB-569 alone, CCI-779 alone, or a dilution series of EKB-569 with fixed doses of CCI-779. Following growth for 3 days, cell survival was determined using the SRB test procedure. Cytotoxicity curves
15 are shown in Fig. 1. EKB-569 produced an IC_{50} value of 0.31 $\mu\text{g/ml}$ in HCT116 cells. When this compound was combined with 2.08 $\mu\text{g/ml}$ CCI-779 (which caused 41% inhibition of growth when administered alone), the IC_{50} value is reduced to 0.03 $\mu\text{g/ml}$, a 10-fold decrease. When combined with 0.026 $\mu\text{g/ml}$ CCI-779 (which alone inhibits cell proliferation by 36 %), the IC_{50} value dropped to 0.051 $\mu\text{g/ml}$, a 6-fold decrease.
20 Similar results were observed when dose-response curves were produced with CCI-779 in the presence of fixed doses of EKB-569. To identify the nature of this drug interaction, isobolograms (at 50% effect level) of the combination of EKB-569 and CCI-779 were generated (Fig. 2). The isobologram was deeply indented with the concave side up, indicating a substantial synergistic interaction between the two
25 drugs. At the most synergistic point, 0.03 $\mu\text{g/ml}$ of EKB-569 combined with 0.077 $\mu\text{g/ml}$ CCI-779 was iso-effective with 0.31 $\mu\text{g/ml}$ of EKB-569 alone or 4.3 $\mu\text{g/ml}$ CCI-779 alone (IC_{50} for each drug alone). Thus, a 10-fold reduction in the dose of EKB-569 and a 50-fold reduction in the dose of CCI-779 was required to inhibit cell proliferation by 50% when the drugs were combined, compared to either drug alone.
30 Isobolograms derived from different endpoints, ranging from 50 to 65% were also examined. As shown in Fig 3., the isobolograms produced were almost superimposable, indicating synergy at all effect levels tested.

The interaction between EKB-569 and CCI-779 was also evaluated using a 3-dimensional analysis. Here, pharmacological interactions are presented in a 3-dimensional plot with the plane at 0% representing additive interaction, and peaks and valleys representing areas of synergy or antagonism, respectively, between the two drugs. In Fig. 4, the combination of EKB-569 and CCI-779 resulted in a broad area of synergistic interaction, consistent with the results shown in the isobologram studies. A contour plot of the 3-dimensional synergy plot facilitates the identification of the concentrations of drugs at which greatest synergistic toxicity occurs (Fig. 5). A broad area of synergy was observed at 0.0005 to 3 $\mu\text{g/ml}$ CCI-779 and 0.16 to 0.4 $\mu\text{g/ml}$ EKB-569. Within this area, two peaks of maximum synergy occurred at 0.0005 to 0.003 $\mu\text{g/ml}$ and 0.05 to 0.3 $\mu\text{g/ml}$ of CCI-779 and 0.25 to 0.37 $\mu\text{g/ml}$ EKB-569.

Based on the results of these standard pharmacological test procedures, combinations of CCI-779 plus EKB-569 acted synergistically together, and are useful as antineoplastic therapy. More particularly, these combinations are useful in the treatment of renal carcinoma, soft tissue sarcoma, breast cancer, neuroendocrine tumor of the lung, cervical cancer, uterine cancer, head and neck cancer, glioma, non-small cell lung cancer, prostate cancer, pancreatic cancer, lymphoma, melanoma, small cell lung cancer, ovarian cancer, colon cancer, esophageal cancer, gastric cancer, leukemia, colorectal cancer, and unknown primary cancer. As these combinations contain at least two active antineoplastic agents, the use of such combinations also provides for the use of combinations of each of the agents in which one or both of the agents is used at subtherapeutically effective dosages, thereby lessening toxicity associated with the individual chemotherapeutic agent.

In providing chemotherapy, multiple agents having different modalities of action are typically used as part of a chemotherapy "cocktail." It is anticipated that the combinations of this invention will be used as part of a chemotherapy cocktail that may contain one or more additional antineoplastic agents depending on the nature of the neoplasia to be treated. For example, this invention also covers the use of the CCI-779/EKB-923 combination used in conjunction with other chemotherapeutic agents, such as antimetabolites (i.e., 5-fluorouracil, floxuradine, thioguanine,

cytarabine, fludarabine, 6-mercaptopurine, methotrexate, gemcitabine, capecitabine, pentostatin, trimetrexate, or cladribine); DNA crosslinking and alkylating agents (i.e., cisplatin, carboplatin, streptazoin, melphalan, chlorambucil, carmustine, methchloroethamine, lomustine, bisulfan, thiotepa, ifofamide, or cyclophosphamide);
5 hormonal agents (i.e., , tamoxifen, roloxifen, toremifene, anastrozole, or letrozole);
antibiotics (i.e., plicamycin, bleomycin, mitoxantrone, idarubicin, dactinomycin, mitomycin, doxorubicin or daunorubicin); immunomodulators (i.e., interferons, IL-2, or BCG); antimitotic agents (i.e., estramustine, paclitaxel, docetaxel, vinblastine, vincristine, or vinorelbine); topoisomerase inhibitors (i.e., topotecan, irinotecan,
10 etoposide, or teniposide.); and other agents (i.e., hydroxyurea, trastuzumab, altretamine, retuximab, L-asparaginase, or gemtuzumab ozogamicin).

As used in this invention, the combination regimen can be given simultaneously or can be given in a staggered regimen, with CCI-779 being given at a
15 different time during the course of chemotherapy than EKB-923. This time differential may range from several minutes, hours, days, weeks, or longer between administration of the two agents. Therefore, the term combination does not necessarily mean administered at the same time or as a unitary dose, but that each of the components are administered during a desired treatment period. The agents may
20 also be administered by different routes. For example, in the combination of CCI-779 plus EKB-569, it is anticipated that the CCI-779 will be administered orally or parenterally, with parenterally being preferred, while the EKB-569 may be administered parenterally, orally, or by other acceptable means. These combination can be administered daily, weekly, or even once monthly. As typical for
25 chemotherapeutic regimens, a course of chemotherapy may be repeated several weeks later, and may follow the same timeframe for administration of the two agents, or may be modified based on patient response.

As typical with chemotherapy, dosage regimens are closely monitored by the
30 treating physician, based on numerous factors including the severity of the disease, response to the disease, any treatment related toxicities, age, health of the patient, and other concomitant disorders or treatments.

Based on the results obtained with the CCI-779 plus EKB-569 combinations, it is projected that the initial i.v. infusion dosage of CCI-779 will be between about 0.1

and 100 mg/m², with between about 2.5 and 70 mg/m² being preferred. It is also preferred that the CCI-779 be administered by i.v., typically over a 30 minute period, and administered about once per week. The initial daily dosages of EKB-569 will be between about 1 and 100 mg, with between 5 and 75 mg being preferred. After one
5 or more treatment cycles, the dosages can be adjusted upwards or downwards depending on the results obtained and the side effects observed.

Oral formulations containing the active compounds of this invention may comprise any conventionally used oral forms, including tablets, capsules, buccal
10 forms, troches, lozenges and oral liquids, suspensions or solutions. Capsules may contain mixtures of the active compound(s) with inert fillers and/or diluents such as the pharmaceutically acceptable starches (e.g. corn, potato or tapioca starch), sugars, artificial sweetening agents, powdered celluloses, such as crystalline and microcrystalline celluloses, flours, gelatins, gums, etc. Useful tablet formulations may
15 be made by conventional compression, wet granulation or dry granulation methods and utilize pharmaceutically acceptable diluents, binding agents, lubricants, disintegrants, surface modifying agents (including surfactants), suspending or stabilizing agents, including, but not limited to, magnesium stearate, stearic acid, talc, sodium lauryl sulfate, microcrystalline cellulose, carboxymethylcellulose calcium,
20 polyvinylpyrrolidone, gelatin, alginic acid, acacia gum, xanthan gum, sodium citrate, complex silicates, calcium carbonate, glycine, dextrin, sucrose, sorbitol, dicalcium phosphate, calcium sulfate, lactose, kaolin, mannitol, sodium chloride, talc, dry starches and powdered sugar. Preferred surface modifying agents include nonionic and anionic surface modifying agents. Representative examples of surface modifying
25 agents include, but are not limited to, poloxamer 188, benzalkonium chloride, calcium stearate, cetostearyl alcohol, cetomacrogol emulsifying wax, sorbitan esters, colloidal silicon dioxide, phosphates, sodium dodecylsulfate, magnesium aluminum silicate, and triethanolamine. Oral formulations herein may utilize standard delay or time release formulations to alter the absorption of the active compound(s). The oral
30 formulation may also consist of administering the active ingredient in water or a fruit juice, containing appropriate solubilizers or emulsifiers as needed.

In some cases it may be desirable to administer the compounds directly to the airways in the form of an aerosol.

The compounds may also be administered parenterally or intraperitoneally. Solutions or suspensions of these active compounds as a free base or pharmacologically acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxy-propylcellulose. Dispersions can also be prepared in
5 glycerol, liquid polyethylene glycols and mixtures thereof in oils. Under ordinary conditions of storage and use, these preparation contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of
10 sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol
15 (e.g., glycerol, propylene glycol and liquid polyethylene glycol), suitable mixtures thereof, and vegetable oils.

For the purposes of this disclosure, transdermal administrations are understood to include all administrations across the surface of the body and the inner linings of bodily passages including epithelial and mucosal tissues. Such
20 administrations may be carried out using the present compounds, or pharmaceutically acceptable salts thereof, in lotions, creams, foams, patches, suspensions, solutions, and suppositories (rectal and vaginal).


Transdermal administration may be accomplished through the use of a transdermal patch containing the active compound and a carrier that is inert to the
25 active compound, is non toxic to the skin, and allows delivery of the agent for systemic absorption into the blood stream via the skin. The carrier may take any number of forms such as creams and ointments, pastes, gels, and occlusive devices. The creams and ointments may be viscous liquid or semisolid emulsions of either the oil-in-water or water-in-oil type. Pastes comprised of absorptive powders dispersed in
30 petroleum or hydrophilic petroleum containing the active ingredient may also be suitable. A variety of occlusive devices may be used to release the active ingredient into the blood stream such as a semi-permeable membrane covering a reservoir containing the active ingredient with or without a carrier, or a matrix containing the active ingredient. Other occlusive devices are known in the literature.

Suppository formulations may be made from traditional materials, including cocoa butter, with or without the addition of waxes to alter the suppository's melting point, and glycerin. Water soluble suppository bases, such as polyethylene glycols of various molecular weights, may also be used.

5

The following provides the preparation of EKB-569 from commercially available starting materials or starting materials that can be made according to available literature procedures.


10 **Preparation of 4-dimethylaminocrotonic acid from TMS-4-bromocrotonate**



211 ml dimethylamine (2M in THF, 0.422 moles) was added drop-wise to a solution of 50 g TMS-4-bromocrotonate (0.211 moles, 75.9% by GC-MS) in 250 ml of THF at 0-50°C under N₂. The reaction mixture was stirred at room temperature for 30 minutes.

15 A white solid by-product was filtered off. 2 ml water was added to the filtrate followed by seeding. The crystals formed were filtered and washed with ether to give 18.3 g (from two crops) off-white solid product. Yield was 67.2% (98% purity by GC-MS, NMR was consistent with the structure).


20 **Preparation of methyl 4-dimethylaminocrotonate from methyl-4-bromocrotonate**



120 ml dimethylamine (2M in THF, 0.24 moles) was added drop-wise to a solution of 20 g methyl 4-bromocrotonate (85% purity, 0.095 moles) in 150 ml of THF at 0-50°C under N₂. The reaction mixture was stirred for 15 minutes at room temperature. TLC

(9:1 CH₂Cl₂: MeOH with few drops of Et₃N) showed residual methyl 4-bromocrotonate. The reaction mixture was heated to 40-45°C for 15 minutes. A white solid by-product was filtered off. The filtrate was evaporated to give a yellow oil (14 g). The yellow oil was dissolved in 100 ml CH₂Cl₂ and washed with H₂O twice. The aqueous layer was back extracted with 100 ml CH₂Cl₂. The CH₂Cl₂ layers were combined, dried over MgSO₄ and filtered. The filtrate was evaporated to give an oil (12 g). Yield was 88%. NMR indicated desired product with trace methyl 4-bromocrotonate.

10 **Preparation of Methyl 4-N,N-dimethylaminocrotonate hydrochloride on large scale:**




A 3L flask was charged with tetrahydrofuran (0.71 kg, 0.80 L). Methyl 4-bromocrotonate (0.20 kg, 0.13 L, d = 1.522 g/mL) was added and rinsed with tetrahydrofuran (0.18 kg, 0.20 L). The solution was stirred and cooled to 0-10°C. An additional funnel was charged with a solution of dimethylamine in tetrahydrofuran and added over (1 h 15 min) keeping the temperature at 0-10°C. The mixture was stirred for a minimum of 30 mins and checked for reaction completion by TLC. The reaction was complete when there is ≤ 2% detectable starting material (methyl 4-bromocrotonate) present. The mixture was filtered cold on a Buchner funnel into a 3 L multi-neck flask, rinsed with pre-chilled (0-10°C) tetrahydrofuran (2 x 0.18 kg, 2 x 0.20 L), and suction maintained until dripping stops. The flask was equipped with an agitator, thermometer, and a setup for vacuum distillation. The solution was concentrated by distillation under a reduced pressure of (125-200 mm Hg) and at a maximum pot temperature of (40°C) to a pot volume of (200 mL). Isopropanol (0.22 kg, 0.28 L) was added and the mixture cooled to 0-10°C. The distillation stillhead was replaced with an addition funnel charged with a solution of HCl in isopropanol, which was added over 45 min until pH of 2.0-3.0 was reached, while maintaining a temperature 0-10°C. The mixture was held for a minimum 30 min, and filtered cold on a Buchner funnel, rinsed with isopropanol (2 x 0.12 kg, 2 x 0.15 L). The filter cake was dammed and

suction maintained until dripping stopped. The product was dried in a vacuum oven at 50°C and 10 mm Hg for 18-20 h.

Preparation of 4-dimethylaminocrotonic acid hydrochloride from methyl


5 4-dimethylaminocrotonate



A NaOH solution (3.35 g in 25 ml H₂O, 0.084 moles) was added drop-wise to a solution of 12 g methyl 4-dimethylaminocrotonate (0.084 moles) in 100 ml MeOH at room temperature. The reaction mixture was heated to 40-45°C for 1 hour then cooled to room temperature. The pH was adjusted to 1-2 with 5 N HCl. The mixture was concentrated to a thick oil which was triturated with dehydrated alcohol to form a solid. The solid by-product was filtered off. The filtrate was evaporated to an oil which was triturated with IPA. Seven (7.0) g of white solid product was obtained. Yield was 50% with the purity 86.3 % by GC-MS.

15

Preparation of 4-N,N-dimethylaminocrotonic acid hydrochloride on large scale




A 2 L multi-neck flask was equipped with agitator, thermometer, addition funnel, and nitrogen protection. The flask was charged with ethanol (0.39 kg, 0.50 L). Methyl 4-N,N-dimethylamino crotonate hydrochloride (0.125 kg) was added and rinsed with

20

ethanol (0.10 kg, 0.125 L). The suspension was stirred and cooled to 0-10°C. The addition funnel was charged with sodium hydroxide (50%) (0.11 kg, 0.072 L, d=1.53 g/mL) and added over 20 min keeping the temperature at 0-10°C. A slight exotherm was observed and the mixture turned yellow. The mixture was stirred for a minimum of 15 min, and then warmed to 18-22°C, and held for a minimum of 4 h. The reaction was checked for completion by TLC. The reaction is complete when there is $\leq 2\%$ detectable starting material (methyl 4-N,N-dimethylaminocrotonate hydrochloride) present. The mixture was cooled to 0-10°C. An addition funnel was charged with a solution of HCl in isopropanol and added over 40 min until pH 2.0-3.0 was attained, while maintaining the pot temperature of 0-10°C. The mixture was stirred for a minimum of 30 min, and filtered cold on a Buchner funnel into a 2 L multi-neck flask, rinsed with cold ethanol (0-10°C) (2 x 0.05 kg, 2 x 0.063 L) with suction maintained until dripping stops. The flask was equipped with an agitator, thermometer, and setup for vacuum distillation. Solvent was removed under a reduced pressure of 50-100 mm Hg and at a maximum pot temperature of (40°C) to a pot volume of 160-180 mL. Isopropanol (0.049 kg, 0.063 L) was added, and the mixture warmed to 35-40°C over 10 min. Acetone (0.10 kg, 0.13 L) was added over 20 min while maintaining the pot temperature at 35-40°C. The mixture was seeded and cooled to ambient temperature 20-25°C, and held there for a minimum of 12-18 h. The mixture was cooled to 0-10°C, held there for a minimum of 1 h. A mixture of isopropanol (0.049 kg, 0.063 L) and acetone (0.10 kg, 0.13 L) was prepared, stirred to homogenize, and cooled to 0-10°C. The mixture was filtered cold on a Buchner funnel, rinsed with isopropanol/acetone (2 x 0.074 kg, 2 x 0.096 L), and the filter cake dammed while maintaining suction until dripping stopped. The product was dried in a vacuum oven at 50°C and 10 mm Hg for 18-20 h.

Preparation of 4-dimethylaminocrotonyl anilide from 4-dimethylaminocrotonic acid hydrochloride



Thionyl chloride (0.36 ml, 0.005 moles) was added drop-wise to a solution of 0.33 g 4-dimethylaminocrotonic acid hydrochloride (0.002 moles) in 15 ml CH_2Cl_2 containing 2 drops of DMF at 0°C under N_2 . The reaction mixture was refluxed for 30 min. Then 0.72 ml aniline (0.008 moles) was added drop-wise to the reaction mixture at 0°C and stirred for 1 hour at room temperature. A solid by-product was filtered. The filtrate was evaporated to give an oil (0.6 g). GC-MS data shows that the oil is 11.7% 4-dimethylaminocrotonic acid hydrochloride and 85% of desired product.

Preparation and isolation of 4-N,N-dimethylaminocrotonoylchloride hydrochloride

A well stirred suspension 4-dimethylaminocrotonic acid hydrochloride (5.0 g, 30 mmol) in cold (0°C) THF (40 mL) and DMF (2 pipet drops) was treated with oxalyl chloride (3.15 mL, 36 mmol). The mixture was stirred at $20\text{--}25^\circ\text{C}$ for 3 h then cooled to 0°C and held for 30 min. The solids were collected on Buchner funnel (under a blanket of nitrogen) and washed with cold (0°C) THF (3 x 5 mL). The product was dried under vacuum (~ 1 torr) at $40\text{--}50^\circ\text{C}$ for 3 h to give 4.0 g of 4-dimethylaminocrotonoyl chloride hydrochloride. This material is characterized as its methyl ester by treatment of the solid with methanol.

Alternatively, the title compound can be prepared in CH_3CN and used directly for the coupling step:

Preparation of EKB-569

A 3 L multi-neck flask was equipped with an agitator, thermometer, dip tube, and nitrogen protection. The flask was charged with N-methyl pyrrolidinone (0.77 kg, 0.75 L, $d=1.033$ g/mL). At ambient temperature, 4-[3-chloro-4-fluorophenyl]amino-6-amino-3-cyano-7-ethoxy quinoline (0.0748 kg) [see, US Patent 6,002,008] was added and the mixture stirred while heating to 40-45 °C and hold for 15 min. The flask was cooled to 0-10°C. The mixture containing 4-N,N-dimethylaminocrotonoyl chloride hydrochloride was transferred via dip tube and positive nitrogen pressure to the 3 L flask over 30-45 min, while maintaining 0-10°C. The mixture was kept at 0-10°C for a minimum of 2 h. The reaction was checked for completion by HPLC. The reaction is complete when there is $\leq 2\%$ of the starting material (4-[3-chloro-4-fluorophenyl]-amino-6-amino-3-cyano-7-ethoxy quinoline) present. A 12 L multi-neck flask equipped with agitator, thermometer, dip tube, and nitrogen protection was charged with water (2.61 kg, 2.61 L). Sodium bicarbonate (0.209 kg) was added and stirred until a solution was obtained. The solution was cooled to 20-24°C. The NMP-CH₃CN mixture was transferred, via dip tube and positive nitrogen pressure, to the 12 L flask over 45-60 min, while maintaining 20-24°C. The mixture was maintained at 20-24°C for a minimum of 1 h, and filtered on a Buchner funnel, and rinsed with water (3 x 0.40 kg, 3 x 0.40 L) with suction being maintained until dripping stops. The product was dried in a vacuum oven at 50°C and 10 mm Hg for 28-30 h to give 78.5 g (86% yield) of product.

CLAIMS

1. A method of treating a neoplasm in a mammal in need thereof, which
5 comprises providing to said mammal an effective amount of a combination comprising
CCI-779 and EKB-569.
2. The method according to claim 1, wherein the neoplasm is renal cancer.
- 10 3. The method according to claim 1, wherein the neoplasm is soft tissue
sarcoma.
4. The method according to claim 1, wherein the neoplasm is breast
cancer.
- 15 5. The method according to claim 1, wherein the neoplasm is a
neuroendocrine tumor of the lung.
6. The method according to claim 1, wherein the neoplasm is cervical
20 cancer.
7. The method according to claim 1, wherein the neoplasm is uterine
cancer.
- 25 8. The method according to claim 1, wherein the neoplasm is a head and
neck cancer.
9. The method according to claim 1, wherein the neoplasm is glioma.
- 30 10. The method according to claim 1, wherein the neoplasm is non-small
cell lung cancer.
11. The method according to claim 1, wherein the neoplasm is prostate
cancer.

12. The method according to claim 1, wherein the neoplasm is pancreatic cancer.
- 5 13. The method according to claim 1, wherein the neoplasm is lymphoma.
14. The method according to claim 1, wherein the neoplasm is melanoma.
- 10 15. The method according to claim 1, wherein the neoplasm is small cell lung cancer.
16. The method according to claim 1, wherein the neoplasm is ovarian cancer.
- 15 17. The method according to claim 1, wherein the neoplasm is colon cancer.
18. The method according to claim 1, wherein the neoplasm is esophageal cancer.
- 20 19. The method according to claim 1, wherein the neoplasm is gastric cancer.
20. The method according to claim 1, wherein the neoplasm is leukemia.
- 25 21. The method according to claim 1, wherein the neoplasm is colorectal cancer.
22. The method according to claim 1, wherein the neoplasm is unknown primary cancer.
- 30 23. A method according to any one of Claims 1 to 22 which comprises providing to said mammal an effective amount of a combination comprising CCI-779

and EKB-569, wherein either CCI-779, EKB-569, or both are provided in subtherapeutically effective amounts.

24. A method according to claim 23 in which CCI-779 is provided in a
5 subtherapeutically effective amount.

25. A method according to claim 23 in which EKB-569 is provided in a subtherapeutically effective amount.

10 26. A method according to claim 23 in which both CCI-779 and EKB-569 are provided in subtherapeutically effective amounts.

27. An antineoplastic combination which comprises an antineoplastic effective amount of a combination of CCI-779 and EKB-569.
15

28. A composition according to claim 27 in which CCI-779 is provided in a subtherapeutically effective amount.

29. A composition according to claim 27 in which EKB-569 is provided in a
20 subtherapeutically effective amount.

30. A composition according to claim 27 in which both CCI-779 and EKB-569 are provided in subtherapeutically effective amounts.

25 31. A product comprising CCI-779 and EKB-569 as a combined preparation for simultaneous, separate or sequential use in the treatment of a neoplasm in a mammal

30 32. A product according to claim 31 in which the neoplasm is one of the following: renal cancer, soft tissue sarcoma, breast cancer, neuroendocrine tumor of the lung, cervical cancer, uterine cancer, a head and neck cancer, glioma, non-small cell lung cancer, prostate cancer, pancreatic cancer, lymphoma, melanoma, small cell lung cancer, ovarian cancer, colon cancer, esophageal cancer, gastric cancer, leukemia, colorectal cancer or unknown primary cancer.

33. Use of CCI-779 and EKB-569 in the preparation of a medicament for the treatment of neoplasm in a mammal.

5 34. A use according to claim 33 in which the neoplasm is one of the following: renal cancer, soft tissue sarcoma, breast cancer, neuroendocrine tumor of the lung, cervical cancer, uterine cancer, a head and neck cancer, glioma, non-small cell lung cancer, prostate cancer, pancreatic cancer, lymphoma, melanoma, small cell lung cancer, ovarian cancer, colon cancer, esophageal cancer, gastric cancer,
10 leukemia, colorectal cancer or unknown primary cancer.

35. A use according to claim 33 or 34 in which CCI-779 is provided in a subtherapeutically effective amount.

15 36. A use according to claim 33 or 34 in which EKB-569 is provided in a subtherapeutically effective amount.

37. A use according to claim 33 or 34 in which both CCI-779 and EKB-569 are provided in subtherapeutically effective amounts.

20

1/5




FIG.1

2/5




FIG.2

3/5




FIG.3

4/5





FIG.4

5/5





FIG. 5

INTERNATIONAL SEARCH REPORT

PCT/US 02/24841

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 A61K31/395 A61P35/00 A61K31/4706 //(A61K31/395,31:4706)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

MEDLINE, EMBASE, EPO-Internal, WPI Data, PAJ, SCISEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	GUPTA R A ET AL: "Combinations for cancer prevention." NATURE MEDICINE, (2000 SEP) 6 (9) 974-5., XP002223840 the whole document, especially from page 974, right-hand column, line 17 to page 975, left-hand column, line 18	1-37
A	TORRANCE C J ET AL: "Combinatorial chemoprevention of intestinal neoplasia." NATURE MEDICINE, (2000 SEP) 6 (9) 1024-8, XP002223841 the whole document -- -/--	1-37



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

9 December 2002

Date of mailing of the international search report

10/01/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+31-70) 340-3016

Authorized officer

Venturini, F

2

Form PCT/ISA/210 (second sheet) (July 1992)

page 1 of 2

INTERNATIONAL SEARCH REPORT

PCT/US 02/24841

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>GREENBERGER L M (REPRINT) ET AL: "EKB - 569: a new irreversible inhibitor of epidermal growth factor receptor tyrosine kinase for the treatment of cancer." CLINICAL CANCER RESEARCH, (NOV 2000) VOL. 6, SUPP. 'S', PP. 388-388. PUBLISHER: AMER ASSOC CANCER RESEARCH, PO BOX 11806, BIRMINGHAM, AL 35202. ISSN: 1078-0432., XP001120789</p> <p>WYETH AYERST RES, PEARL RIVER, NY 10965 abstract</p>	1-37
A	<p>GEOERGER B ET AL: "ANTITUMOR ACTIVITY OF THE RAPAMYCIN ANALOG CCI-779 IN HUMAN PRIMITIVE NEUROECTODERMAL TUMOR/MEDULLOBLASTOMA MODELS AS SINGLE AGENT AND IN COMBINATION CHEMOTHERAPY" CANCER RESEARCH, AMERICAN ASSOCIATION FOR CANCER RESEARCH, BALTIMORE, MD, US, vol. 4, no. 61, 15 February 2001 (2001-02-15), pages 1527-1532, XP001079526</p> <p>ISSN: 0008-5472</p> <p>the whole document</p>	1-37
A	<p>US 5 066 493 A (SEHGAL SURENDRA N ET AL) 19 November 1991 (1991-11-19)</p> <p>the whole document</p>	1-37
P,A	<p>WO 02 13802 A (AMERICAN HOME PROD) 21 February 2002 (2002-02-21)</p> <p>the whole document</p>	1-37

INTERNATIONAL SEARCH REPORT

PCT/US 02/24841

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 1-26 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

PCT/US 02/24841

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5066493	A	19-11-1991	US 5206018 A	27-04-1993
			US 4885171 A	05-12-1989
			BE 877700 A1	14-01-1980
			JP 55073616 A	03-06-1980
			ZA 7905449 A	26-11-1980
WO 0213802	A	21-02-2002	AU 8313901 A	25-02-2002
			WO 0213802 A2	21-02-2002
			US 2002045638 A1	18-04-2002

Form PCT/ISA/210 (patent family annex) (July 1992)

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)